



Mapping the Expertise and Understanding of Menarche, Menstrual Hygiene, and Menstrual Health among Adolescent Ladies in Low- and Center-Profit Nations

Rehan Haider

Riggs Pharmaceuticals Karachi

Corresponding Author: Rehan Haider rehan_haider64@yahoo.com

ARTICLE INFO

Keywords: Adolescent Girls, Menarche, Menstrual Health, Menstrual Health Problems, Menstrual Hygiene Management, Menstruation

Received : 17 May

Revised : 19 June

Accepted: 21 July

©2023 Haider: This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/).



ABSTRACT

Many adolescent girls begin their periods uninformed and unprepared; moms are the primary supply of facts, however, they tell girls, too-little and too past due and regularly speak their misconceptions and due to the fact menstruation is widely visible as polluting and shameful, girls are often excluded and shamed in their houses and their groups. Many no longer have the manner for self-care and no longer get the help they need when they face issues, which averts their potential to hold on with regular activities and can also set up a basis for lifestyles-long dis empowerment. Efforts to respond to girls' needs have been fragmented and piecemeal. There is a growing acknowledgment that efforts are much more likely to be successful if they come together in an entire-of-community technique that includes schools, fitness centers, and houses and communities; educate girls about menstruation, Create norms that see menstruation as healthful and nice, no longer shameful and dirty, enhance get admission to sanitary products, strolling water, and functional bathrooms and privateness for self-care, enhance care for and aid with the aid of ladies' families after they have their periods, enhance access to in a position and caring medical experts once they experience menstrual health troubles

INTRODUCTION

Girls in many low- and middle-earnings worldwide locations (LMIC) enter puberty with facts gaps and misconceptions about menstruation, unprepared to manage it, and uncertain of whilst and in which to look for assistance. This is because adults around them, together with dad, mom, and teachers, are themselves sick-knowledgeable and uncomfortable discussing sexuality, reproduction, and menstruation (which regularly comes weighted down with dirty, polluting, and shameful connotations). To respond to the multiplied worldwide attention on empowering women through the United Nations' Sustainable Improvement Desires, targets were compared to map the knowledge, attitudes, ideals, and practices surrounding menarche, menstrual hygiene, and menstrual fitness among adolescent women in LMIC to inform the future design of applicable suggestions and programming. To do Our dreams are to answer the following questions: (1) understanding and knowledge are girls in LMIC about menstruation and the way they are organized to reach menarche; (2) who are their assets of information regarding menstruation; (3) understand properly the adult's spherical them reply to their information needs; (4) what terrible fitness and social effects do children enjoy due to menstruation and (five) information do teenagers reply after they enjoy the terrible effects and what practices do they expand as a result?

LITERATURE REVIEW

Whether by way of a relative, friend, or some other community member, the statistics on menstrual health and hygiene provided by the two teens were not constantly well-timed, nor was it good enough.

Researchers have determined that mothers, spouses, and children in India [1, 7] and Tanzania [49], who offered girls' information, often did not achieve this until after menarche. In Mexico, however, 94.0% of girls stated that they had discussed menstruation with their mothers before menarche [24]. A look at Nigeria stated that 55.2% of faculty-going girls have been "trained" before attaining menarche, which protected them from being made aware of what to count on at menarche and how to accumulate blood and do away with substances [40]. Similarly, dad and mom's training degree was found to have an extensive effect on pre-menarcheal knowledge in Nigeria ($p < \text{zero}.05$) as women whose dad and mom had acquired tertiary schooling were most likely to have been killed [40]. In India [15, 33] and Kenya [50, 51], women stated that few facts about menstruation changed into furnished and almost no rationalization. assets of information may additionally have misconceptions about menstruation, which they'll skip on. mothers interviewed in Bangladesh attributed menstruation to God [28]. all through initiation rites in Malawi, misconceptions, like men can get hurt if they come in contact with menstrual blood and are instructed to girls via a lady household [52].

Given the hyperlink between menstruation and the potential to conceive, mothers interviewed in Bangladesh did not forget it appropriate to speak about the count number with their pre-menarcheal daughters [28]. Each mother and instructor, most of whom were male, incited pain as an obstacle to discussing menstruation with ladies [53]. Instructors in rural Tanzania warned their college

students that their mothers might be disillusioned if they were instructed to accomplish menarche [49]. This could be a result of the cultural taboos that prevented parents from discussing sex-related subjects with their daughters. Taboos were also noted with the aid of a few instructors in Tanzania who desired to offer support to their college students [54]. Instructors in Kenya do not perceive menstrual education as important as part of their function, nor did they sense well prepared to share statistics for the students [53].

A majority of teachers (70–90%) at colleges in Ghana who had been trained The application of play-based total procedures to sell menstrual understanding and practices had been confident in discussing menstruation with their college students as compared to their opposite numbers at faculties, not in the usage of comparable approaches that had restricted conversations [55]. standard, 82.4.% of have a look at participants in Jordan felt they had been not thoroughly prepared for reaching menarche [36]. Among girls in rural Nepal who obtained statistics from a parent, pal, or coursebook, an overwhelming majority felt that menstrual-related topics were no longer well taught [5]. 4-5th through fifth faculty-going women in Egypt wanted more information [30]. However, girls in Malaysia [39] and Tanzania [54] expressed feelings.ashamed, embarrassed, and uncomfortable when inquiring about menstruation in adults.

What are the emotional, physical, and social effects of menstruation on girls?

The expected and experienced impacts related to menstruation were mostly negative (with some positive impacts when girls were better informed/prepared). While eight out of ten study participants in Mexico expected at least one positive change after reaching menarche, all expected to say that the least one negative change occurred, and the nine most anticipated changes between cities and rural girls were negative, such as discomfort, apprehension, and cramps [56]. Overall, 89.4% of expected changes reported by premenarcheal girls and 88.7% of changes in girls after menarche were negative [57]

Emotional Impacts

A quarter to 8 out of 10 girls questioned in numerous LMICs is now not mentally organized to reach menarche (Brazil [58], China [4], India [11], Jordan [36], Mexico [24, 56], Nepal [25], Nigeria [40],). Many women experienced terrible reactions during the first period (Table 46.4). For instance, a majority of college-going women in a single country take a look at in India defined menarche as a shocking or apprehensive event and many cried upon seeing their blood [18]. A few faculty-going women perceived menstrual blood as dirty or described feeling disgusted by their length (30. 5% in Lebanon [59], and 48. 9%, as opposed to 72. .8% in rural areas as opposed to Malaysia [38, 48], and 10.0–23.4% in Indian states [11, 17, 34]. additionally, girls in Kenya found out that “the girl along with her length is the only one to hang her head” because she becomes the target of undesirable and on occasion unkind interest [53]. temper swings and irritability linked to menstruation were each suggested by extra than -thirds of schoolgirls in India [60], Lebanon [59], and Malaysia [38, 48]. Currently, not all emotions

about achieving menarche have been terrible; more than 1/2 of schoolgirls in China [4], India [9], and Malaysia [38, 48] felt pleasure in maturing. Attention organizations in rural and urban settings in Kenya [51] and Tanzania [61] with female college students have discovered similar sentiments. The more college-going women in Mexico knew about menstruation, the less negative their attitudes ($p < \text{zero}.05$); and the greater the organization they felt, the more superb their attitudes ($p < 0.0001$) [24]. A later menarcheal age and higher socioeconomic status appeared to reduce the incidence of adverse reactions. Amongst girls in Malaysia [39] and Turkey [2], respectively

Table 1. Negative Reaction Upon Reaching Menarche

<i>First author, Year</i>	<i>Country</i>	<i>Setting</i>	<i>School status</i>	<i>N</i>	<i>Negative reaction^a (%)</i>
West Africa					
Aniebue 2009 [40]	Nigeria	Urban	School-going	495	50.3%
Oche 2012 [14]	Nigeria	Urban	School-going	122	53.3%
Central America					
Marván 2001 [57]	Mexico	Urban	School-going	98	15.3%
East Asia					
Tang 2003 [4]	China	Unclear	School-going	1,573	72.0%
South Asia					
Bosch 2008 [28]	Bangladesh	Rural	Unclear	86	64.0%
Mudey 2010 [34]	India	Rural	School-going	300	43.7%
Shanbhag 2012 [10]	India	Rural	School-going	329	44.1%
Tiwari 2006 [11]	India	Unclear	School-going	763	20.6%
Bobhate 2011 [7]	India	Urban	Unclear	241	64.7%
Udgiri 2010 [23]	India	Urban	Unclear	342	31.0%
Adhikari 2007 [5]	Nepal	Rural	School-going	150	96.7%
Ali 2010 [6]	Pakistan	Urban	Government school	425	55.8%
Ali 2010 [6]	Pakistan	Urban	Out-of-school	425	62.6%
Ali 2010 [6]	Pakistan	Urban	Private school	425	55.1%
West Asia					
Reis 2011 [27]	Turkey	Urban	Mix	310	43.9%
Ersoy 2004 [2]	Turkey	Urban	School-going	1,017	49.8%

Physical Impacts

The physical effects of menstruation that were generally reported throughout studies blanketed premenstrual signs, symptoms, syndromes, and painful intervals. These outcomes were almost consistently suggested as a minimum of half of the pattern (Table 46.5). A maximum of 93.2% of rural women in India reported experiencing premenstrual symptoms [62] and 94.4% of school-going women in Egypt reported experiencing dysmenorrhea [44]. In Ethiopia, women with premenstrual symptoms experienced dysmenorrhea more often than those without (82.4% vs. 40. %, respectively) [21]. Of ladies with dysmenorrhea in

In Ghana, nearly two-thirds of patients experience symptoms throughout the maximum number of cycles [31]. A majority of rural-living women surveyed in Malaysia considered dysmenorrhea a normal issue of menstruation [47]

Social Impacts

A quarter of the women in rural India [63], a third of the female students, had daily sports or daily exercise restricted by menstruation in Brazil [58] and Egypt [44], and 60.0% of slum dwellers in India [64]. In urban Sri Lanka, schoolgirls with PMS had significantly more disruption in daily exercise than those who did not. [43]. Every day, sports are similarly restricted by taboos related to what and who men chase women can come into contact with. Menstruating women in India and Nepal are now prevented from entering

kitchens or bedrooms, ensuring that menstrual blood does not contaminate food or other sources [1, 5, 11, 12,63, 65]. Household chores that include cooking are regularly marked as “not now allowed” for menstruating women in India [1, 12, 63], Kenya [53], and Nepal [5]. female university students from a mix of rural and urban settings in India reported restrictions on who they can contact at the same time as they are menstruating [12, 18]. Other frequently reported social barriers include the avoidance of physical or social activities(e.g., sports activities and abilities) and refraining from or missing spiritual activities faculties (Table 46.6). Women in Malaysia [38, 48] and Pakistan [6] have suggested limiting their religious activities because of menstruation. studies reporting complete abstinence from religious sports typically come from India; this exercise turned into reported with the help of 44.7-94.2% of the interviewed ladies [7-10, 12, 13, 15, 34, 60, 63,64]. level of training had a significant effect on diversion practice in holy places in India ($p < 0.05$) [23]. Mothers interviewed in Nigeria found that they advised their daughters to refrain from praying during their period [41]. This is in line with all the others, look at Nigeria where there are 43. of women declared that they abstain from religious activities [14]. fitness exercise interventions in Bangladesh [66] and India [19] did not result in significant a decline in religious restrictions among women throughout their duration. even when asked if girls can transfer to high school even if they are menstruating, 70.7% of girls in rural Nepal answered “no” [5]. actual absence mentioned in various LMICs has not yet reached this level, as a replacement starting at 2.0% of the urban population Womens in Nigeria [40] to 61.7% of women living in rural Uganda [16]. awareness agencies in Malaysia revealed that dysmenorrhea may also have a greater effect on the absence of women in faculties in the city than in the countryside[38]. Dysmenorrhea turned into significantly associated with missing teachers girls living in a city in Lebanon [59] became a serious pain among those in Brazil [58] ($p\text{-fee} < \text{zero}.05$). Menstrual disorders have generally become markedly associated with a lack of college education among women living in rural India ($p\text{-price} < \text{zero}.001$) [67]. In Kenya, male instructors reportedly teased girls about menstruation after they disappeared after a few days of secondary school [53]. Although the teachers denied this, they mentioned that they were involved in maidens that were scattered in magnificence [53]. said that teachers interviewed in Ghana have comparable problems with ladies who are scattered and out of school [68]. Girls in India [60], Malaysia [47, 48], and Uganda [16] related menstruation to negative overall academic performance and intermittent grades.

A puberty schooling intervention with the provision of sanitary pads in a non-randomized trial in Ghana significantly improved women’s school attendance ($p\text{-Value} < 0.001$) [69]. As an alternative, a randomized trial in Nepal [70] tested that presenting menstrual cups may additionally enhance convenience and mobility, and one in Kenya [71] established that they can reduce distractions related to leakage and improve college attendance.

How do women cope with dire consequences, and What Practices Do They expand? To cope with the physical influences of menstruation described in Table 46.5, some girls reported the use of traditional remedies (Bangladesh [72], Brazil [58], India [63, 64, 73, 74], Malaysia [37, 47, 48, 75], Sri.Lanka [43],) and others stated taking medicinal drugs to alleviate aches, frequently utilizing manner of self-medicating or consulting pharmacies (Bangladesh [72], Brazil[58], Egypt [44], India [60, 63, 64, 74], Iran [76], Malaysia [39, 47, 75], Nigeria [42], and Turkey [27, 77]). consultation of fitness specialists for menstrual-associated troubles became minimum, usually cited by using much less than a fifth of girls (Bangladesh [72], Brazil [58], Ethiopia [21], India [60, 63,64, 73, 74], Iran [76], and Malaysia [37, 47, 48],). but, one examiner did file that 69.eight% of Indian girls with issues sought interest from a health expert [7]. some others take a look at India and said that 19.2% of girls with a hassle in no way discussed it with all people—a fitness expert, relative, or pal [64]. Girls in Bangladesh were significantly more likely to seek advice from a person regarding their problems after participating in 12 fitness training periods over the route of six months than at baseline ($p < 0.01$) [66]. fashionable lack of person steering related to menstruation may also contribute to the version of simple hygiene management practices that incorporate the use of sanitary absorbents and daily bathing throughout the LMIC (Table 46.7). The use of sanitary pads to absorb blood ranged from 2.zero% of schoolgirls in rural Nepal [5] to 69.1- Approximately 93.8% of urban women live in Nigeria [14, 20, 40, 41]. However, one study of women in rural areas of seven Indian states reported greater proportions of women using material in comparison to sanitary pads [1, 8, 10, 12, 15, 18, 29, 34, 63, 67, 78]. Sanitary pad use has become significantly better among urban girls in India [22] and Ethiopia [21], as modified into the use of sanitary pads or new fabric among school-going women in India [1] ($p < zero.01$). This quasi-experimental study examined village-based conferences for women in India as a dissemination platform.

Health messages have contributed to a significant boom in the use of sanitary pads and a decrease in the reuse of cloth ($p < 0.05$) [3]. At the same time, fewer women using sanitary pads in an unmarried state take a look at pronounced horrific ft and rashes in rural India than did women using material [63], and the value of sanitary pads has been a challenge for some ladies puzzled in one of the studies in India [12, 15, 34], Tanzania [61], and Uganda [16]. Nearly all 102 urban-residing girls puzzled in Kenya desired sanitary pads for their convenience and reliability, but almost 1/2 used fabric or an aggregate of fabric and sanitary pads to save money [50]. Tissue paper and cotton were also mentioned as absorbents for girls in diverse LMICs, with tissue paper being mentioned as utilizing as many as 37.1% of rural-residing faculty women in Uganda [16]. A minority of women in Egypt [30] and India [12, 46], no more than one in five, compared to 56. Five% of women in Nigeria [20] changed their absorbents, even within the faculty. Most women in Egypt [30] and Uganda [16] felt that colleges lacked privacy, and most women in India [12] and Nigeria[20] favored the exchange of materials at home. Insufficient latrines, water materials, and disposal infrastructure provide a barrier for college children in India [15, 67],

Tanzania [61], and Uganda [16] to manipulate their periods at faculty. Methods for removing substances beyond throwing them away with different trash-blanketed burning, burying, and flushing substances. very few (2.5%) girls in Egypt do away with absorbents using burning them in comparison to 17.0–76.0% in India [12, 63, 73] and Nigeria [14, 40]. Indian girls in rural settings were significantly more likely to file burning materials than those in urban settings ($p < \text{zero}.05$) [22]. Among people who reuse materials, drying washed materials in daylight in place of hiding varies from 30.7% of women in urban Pakistan [6] to 44.3–72.4% of schoolgirls in India [10, 12, 15, 18, 73]. School-based health training in India has resulted in significant improvement.

Washing materials with cleaning soap, drying them in the sun, and getting rid of them accurately [17]. pronounced bathing practices in India ranged from all 2 hundred rural-living women in one kingdom abstaining in the course of menstruation [29] to nearly all three,443 women in urban regions in another state bathing each day [73]. One study discovered that the exercise of daily bathing changed significantly more among city-living women than among rural-residing women ($p < 0.05$) [65], and others discovered that each regular supply of water and a personal rest room, one of a kind to a family, had significant relationships with taking each day tub ($p < \text{zero}.001$) [13]. In Turkey [27] and Nigeria [20], eleven.9% and 72.5% of city-living girls, respectively, suggested a growing range of baths. Girls in rural Kenya found out that they wanted to shower more often throughout their period; however, they were worried about the usage of constrained water resources and feared revealing to their circle of relatives that they had been menstruating [51]. A quasi-experimental study of 698 women in Iran showed that individuals who participated in 10 two-hour instructor-led sessions on pubertal changes engaged in normal bathing more than individuals who did not, and the difference was significant ($p < \text{zero}.01$) [79]. Other quasi-experimental examinations in Egypt discovered a significant boom in the variety of women bathing each day after participating in four 30–45 main fitness education periods centered on menstruation ($p\text{-cost} < \text{zero}.001$) [80]. Four studies in India defined pleasant cleansing of genitalia as washing two or more instances a day while menstruating; one-1/3 to 3-quarters of girls met this criterion [12, 13, 29, 34]. Unsatisfactory cleaning is significantly higher among rural living women than among urban residents [22]. In a study evaluating the impact of college-based health training, the share of women using cleaning soap to ease their genitalia significantly multiplied from 30. zero to 94.3% ($p\text{-cost} < \text{zero}.01$) [17]., Girls in Mexico who had previous knowledge of the physiology of menstruation were significantly more likely to know what was happening in their bodies and what to do, in terms of hygiene management, upon attaining menarche ($p\text{-price} < \text{zero}.001$) [24]. Schoolgirls were also prepared in Jordan previous to menarche led to more superb attitudes, and mindset had a significant superb correlation with practices ($p\text{-value} < 0.05$) [36]. training status among women in Iran also had a significant correlation with practice ($P < 0.01$) [6,66].

METHODOLOGY

Our literature seeks to become aware of articles that evaluated the understanding of women regarding menstruation, their record sources, the health and social consequences of menstruation, and the way children and adults responded to these results. We searched the Google Pupil, PubMed, and EBSCO global fitness databases for articles in peer-reviewed journals published between 2000 and 2015. To understand the relevant literature, we used the following search method, (menarche or menstruation or menstrual fitness or menstrual hygiene or guys walk management) and (early life or adolescent or kids or younger) and (female, or female), and (knowledge, perception, exercise, or experience). Via an identified and abstract overview, papers in English that addressed the stories of adolescent girls (aged 10–19 years) in LMIC were retained. Complete-text articles were reviewed to determine whether they addressed one or more of these five questions. Given the limited research to be conducted, descriptive overviews, interventions, and the use of quantitative, qualitative, or mixed strategies of any sample length were all protected. 2 while the focal point of this paper is menstrual reports. Research on the preparedness and attitudes of pre-menarcheal girls has been protected as long as data were stratified through individuals who had and had not reached menarche. To complement our search, we reviewed the reference lists of the included articles and identified a small variety of extra studies that met those huge criteria. S ubsequently, we searched and covered guides via United International locations organizations and global non-governmental groups that answered how groups and their LMIC partners respond to the wishes of girls and make remarks on grammar, punctuation, vocabulary, and sentence shape.

RESULTS AND DISCUSSION

A complete of 81 articles have been identified after discarding reproduction articles and people who did not meet the inclusion standards.

- How knowledgeable are women regarding Menstruation?
- How prepared were they to accomplish menarche?

Women throughout LMIC have limited know-how and knowledge of menstruation before reaching menarche. The proportion of women that were consciously ranged from 2.8% of rural ladies puzzled in Rajasthan, India [1] to for all urban women in Turkey [2] (table 46.2). Village-based meetings for women in Maharashtra, India tested as a platform for disseminating fitness messages and notably contributed to growth from 35.1% of girls interviewed in 2003 to fifty-five. Four% of the women interviewed in 2007 were aware of menstruation earlier than its onset ($p < \text{zero}.05$) [3].

Three-quarters of the 1,573 Chinese women surveyed rated their menstrual understanding as inadequate or insufficient [4]. Again, girls with any information frequently maintained misconceptions about menstruation. For example, a study conducted in rural Nepal reported that 6. Of the and fifty girls surveyed recognized menstruation as a physiological technique while eighty-two. Zero% believed it

Table 2. Characteristics

Study ics	Characteristic	Frequency
	Design	
	Descriptive	70
	Intervention	11
	Method	
	Mixed methods	16
	Qualitative	7
	Quantitative	58
	Region	
	East Africa (Ethiopia, Kenya, Malawi, Tanzania, Uganda)	10
	North Africa (Egypt)	4
	West Africa (Ghana, Nigeria)	10
	North/Central America (Mexico)	3
	South America (Brazil)	1
	East Asia (China)	1
	Southeast Asia (Malaysia)	6
	South Asia (Bangladesh, India, Nepal, Pakistan, Sri Lanka)	39
	West Asia (Iran, Jordan, Lebanon, Turkey)	7
	Setting ^a	
	Mix	21
	Rural	23
	Urban	30
	School status ^a	
	Mix of school-going and out-of-school	12
	School-going	63

Changed into a curse [5]. Reported that menstruation is a herbal physical function that is higher at 19.3% in Pakistan [6] and 18. Three–86.3% in five Indian states [1, 7–13] and 96.7% in Nigeria [14]. Menstruation has changed into a curse, disorder, or representation of sin by some women in the Indian states [7–12, 15] and Uganda [16]. Before receiving fitness schooling at the faculty level, and 72. Four% of women in India have considered menstrual blood impure [17]. An extra know-how hole among women is the lack of expertise regarding the origins of menstrual blood; no greater than a 3rd of ladies efficiently identified the uterus as the source of menstrual blood in four Indian states [7, 12, 15, 18, 19] and rural Nepal [5]. An examination in the fourth Indian state reported that almost no women were aware of the source of their blood (2.5%) [8], and at the same time, nearly one-third were aware (63.3%) [13]. In cities in Pakistan [6] and Nigeria [14], 37.2 and seventy-eight.7% diagnosed the uterus because of the source, respectively, compared with 82.9% of college-going girls in rural Uganda [16]. The simplest one-third of rural-residing, high-faculty women surveyed in India associated menarche with the capacity to conceive [10].

Age had a giant effect on slum dwellers' expertise in India, with Older women were more knowledgeable about menstruation than their younger counterparts ($p < 0.05$) [7]. Similar findings were reported among Nigerian schoolgirls ($p < 0.05$) [20]. Compared to those currently not attending school, the focus has shifted to school girls in India [1] and Pakistan [6]. The educational

level had a significant effect on menstrual expertise in India [7] and Nigeria [14] (p-Value 0.05).

Who are the statistical assets of women?in the path of LMIC research, mothers had been frequently the most usually referred to as the source of statistics and recommendations for girls concerning menstruation (table 46. three). In an assessment of ladies living in town additives of Ethiopia [21] and India [22],

Those in rural settings stated that their mothers delivered an awful lot plenty much less frequently (probably because there had been first-rate female partners and kids to whom they would flip). The following mothers were the most unusually useful resources in four Indian states [1, 11, 12, 23]: Mexico [24], Nepal [25], Nigeria [20, 26], Pakistan [6], and Turkey [27], even though they have been utilized by many good deals, much less than a quarter of women. In some contexts, sisters and buddies surpassed mothers due to the truth of the primary delivery of statistics [6, 13, 28, 29].

A majority of research that examined the jobs of instructors and/or fitness professionals as carriers of menstrual information ranked them due to the fact the least common belongings as compared to women's families and pals (Egypt [30], Ghana [31], India [1, 7, 11, 12, 15, 32-35], Jordan [36], Malaysia [37-39], Nepal [25], Nigeria [14, 20, 26, 40-42], Sri Lanka [43], and Turkey [27].)

Instructors have been defined as deliveries with the resource of lot usage by less than 5% of women wandering in three Indian states [1, 32, 33], Nepal [25], and Sri Lanka [43]. At the maximum, one-third of the topics in Nigeria noted teachers as a supply [41]. University college students in Malaysia are much more likely to quote teachers than those in rural settings.

As a delivery, an extremely good range did not encounter menstrual-associated subjects in the university [39]. Hundred tons of less than 1.0% of women in a rural additive in India [33] and city additives in Jordan [36] counseled having acquired records concerning menstruation from fitness professionals. At maximum, 1 / 4 of the test humans in a metropolis in Nigeria stated health experts as a source [41].

Some research endorsed mass media, alongside radio, television, newspapers, magazines, books, and the internet each due to the reality of the satisfactory beneficial, useful resource to be had through girls or as nutritional supplements to specific property of statistics (Egypt [30,44, 45], Ghana [31], India [15, 29, 32, 34, 35, 46], Jordan [36], Malaysia[37-39, 47, 48], Nepal [25], Nigeria [40-42], Sri Lanka [43], and Turkey[2, 27]). At some times, this property was stated by way of greater than 1 / 4 of girls: seventy-. Four% in Nigeria [41], 90 .2% in Egypt [30], and 29.2-40 3.6% in Malaysia [37, 38, 47, 48].women stated not have obtained statistics from any transport in a few studies. As few as 6.8 and seven.0% of ladies in city Nigeria [42] and Egypt [45],respectively, and seven.8% of the mixture of town- and rural-living women in Ethiopia [21] advocated having no delivery. Check that the town- and rural-dwelling girls in India stated 1 / 4 without supply [32]. In rural Nepal, seventy-six. None of the women recommended delivery of menstrual information deliver [5].

CONCLUSIONS AND RECOMMENDATIONS

A vital dilemma of this assessment is that vague measures are frequently used to explain the menstrual reports of girls, which obstructed statistical aggregation and direct comparisons. For example, the research used extraordinary yardsticks for adequate or insufficient knowledge and used the phrases premenstrual syndrome and dysmenorrhea loosely. Additionally, many studies had small sample sizes and relied closely on self-documentation. Few studies have reported low response rates due to pain or barriers in discussing menstruation. Another limitation is that most relevant records from protected studies come from a constrained number of nations and are not consultants for all LMICs. Among the countries covered, LMIC is a top-notch cultural variant, and the effects presented here should be considered in light of these particular perspectives. Despite these obstacles, the proof offered allows for compliance with the following conclusions:

- Vast numbers of ladies in many countries have knowledge gaps and Misconceptions of menstruation. This leaves them unprepared for use. They experienced menarche and reasons for fear and anxiety.
- Mothers, other lady households, and girl peers are fundamental resources for menstruation data and recommendations. The facts they acquire, but, aren't continually timely or good enough. Only a few have to get admission to additional data from assets inclusive of mass media and the Internet.
- Ladies revel in a spread of symptoms in the course of menstruation – aches, Complications and fatigue. These signs, combined with taboos, result in their inability to participate in family, faculty, or social sports.
- Only a few women try to find healthcare after experiencing menstrual fitness problems. Whatever, they will be hotels for family treatment.
- Women in negative urban and rural communities of LMIC are much less probable to obtain and use sanitary pads. Alternatively, they use materials made at domestic with scraps of antique cloth, cotton, paper, and so forth. Loss of privacy, access to clean water, and purposeful toilets make it tougher for them to manage their periods. it is clear that some distance too many women throughout LMIC are struggling with a nearly complete lack of know-how of their everyday organic maturation and its effects, and once they do receive training, nevertheless struggle with insufficient sanitary substances and insufficient physical and emotional support. even though there's no convincing evidence that negative menstrual hygiene management results in ascending reproductive tract infections [81] or reasons for lasting sequelae, this assessment underscores that coming to terms with menarche and navigating the disgrace and sensible challenges related to its management can also purpose women high-quality tension and disappointment. There remains a want for further research into the physical, intellectual, and social effects of such distress. for example, being unprepared for menarche, being excluded and shamed throughout monthly intervals, being hindered in self-care and un cared for when unwell, undermines a woman's experience of being in charge of her life, her feelings of self-worth and her feeling that the people and institutions surrounding her were conscious of

her wishes. The Massive and lasting effects on the lives of women continue to be studied.

In the short term, but, there are intervention studies that reveal Capacity to enhance menstrual knowledge and hygiene control health education interventions like college-primarily based periods examined in India have led to improved information submit-intervention [17], and similar programs in Egypt [80] and Iran [79] have improved the washing practices of ladies all through their intervals. additionally, a quasi-experimental have a look at in India that involved schooling of scientific officers and imparting reference tools brought about statistically significant enhancements in their case management of menstrual fitness issues for female sufferers over a long time of 15 and 24 [82]. a few thrilling initiatives led through academics, worldwide agencies and the private quarter also are beneath manner. Teaching and inspiring mothers and fathers to communicate with their daughters and sons approximately puberty and menstruation is being carried out through the family's reliance on application [83]. A five-12 monthly initiative using Columbia University launched domestically designed puberty booklets for women and boys in Tanzania, Ghana, Ethiopia, and Cambodia; these were embraced by the Ministries of Education and Health in all four countries [84]. Store the youngsters have also evolved workbooks, modeled on those using Columbia College, for girls and boys in Nepal, Uganda, and Malawi and are wearing out puberty training applications in more than one nation [85]. Similar efforts, along with Cycle-Smart [86] and Grow-up Clever [87], are being implemented with the aid of Georgetown College's Institute of Reproductive Health in Rwanda and Guatemala. In 2014, UNESCO published a coverage booklet with steerage to enhance the talent of school administrators and instructors to teach and guide women and boys in school rooms [88]. Procter and Gamble, major manufacturers of sanitary merchandise, have launched verbal exchange programs in numerous global places where they sell merchandise with advertising and advertising strategies aimed at legitimizing their own family speaks of menstruation and attracting and educating girls at the same time as building their self-esteem [89]. At some unspecified times during the Celebrating Womanhood occasion. 2013, the Water Supply & Sanitation Collaborative Council defined menstrual hygiene as an issue and mentioned a three-pronged method that includes breaking the silence around this trouble remember, making sure of hygienic control, and figuring out mechanisms for the relaxed reuse and disposal of materials [90]. Linking menstruation education with efforts to enhance water, sanitation, and disposal facilities in faculties has been actively promoted and completed by UNICEF in the USA and internationally ranges [91]. At the same time, as obligations are important and promising first steps, greater uptake, and determination are needed to fulfill the rights of women associated with menstrual knowledge, fitness, and hygiene. Concerted multi-degree efforts are required to attain this. On the man or woman diploma, women and boys want to be knowledgeable about approximate puberty. On the circle of relatives degree, ladies want steerage throughout their menstrual cycles. At the community

diploma, we need to enhance access to sanitary products in floodwaters, realistic bathrooms, and privacy. We want success.

And worried healthcare folks who can respond to ladies' questions and problems, and provide care once they have menstrual fitness issues. Sooner or later, we need leaders who can trade the perception of menarche and menstruation to surely certainly one of normalcy and promise in choice to disgrace.³

ACKNOWLEDGMENT

Contributions and support of many individuals and organizations. We are deeply grateful to all those who played a role in the success of this project I would like to thank My Mentor [Dr. Naweed Imam Syed Prof Department of Cell Biology at the University of Calgary and Dr Sadaf Ahmed Head of psychophysiology Research Lab University of Karachi their invaluable input and support throughout the research process. Their insights and expertise were instrumental in shaping the direction of this project.

Declaration of Interest I at this moment declare that :

I have no pecuniary or other personal interest, direct or indirect, in any matter that raises or may raise a conflict with my duties as a manager of my office Management, Conflicts of Interest, The authors declare that they have no conflicts of interest, Financial support and sponsorship, Nil.

REFERENCES

- Khanna A, Goyal RS, Bhawsar R. Menstrual practices and reproductive problems Study of adolescent girls in Rajasthan. *J Health Manag.* 2005;7(1):91–107.
- Ersoy B, et al. Effects of different socioeconomic conditions on Menarche in Female Turkish Students. *Early Hum Dev.* 2004;76(2):115–25.
- Dongre AR, Deshmukh PR, Garg BS. The effect of community-based health education interventions on menstrual hygiene management among rural Indian adolescent girls. *World Health Popul.* 2007;9(3):48–54.
- Tang CS, Yeung DY, Lee AM. Psychosocial correlates of emotional Responses to menarche among Chinese adolescent girls. *J Adolescent Health.* 2003;33(3):193–201.
- Adhikari P, et al. Knowledge and practice regarding menstrual hygiene in rural adolescent girls in Nepal. *Kathmandu Univ Med J (KUMJ).* 2007;5(3):382–6
- Ali TS, Rizvi SN. Menstrual knowledge and practices of female adolescents in urban Karachi, Pakistan. *J Adolescent.* 2010;33(4):531–41.
- Bobhate P, Shrivastava S et al.. This was a cross-sectional study of knowledge and practices regarding reproductive health among female adolescents in an urbsluminf Mumbai. *J Fam Reprod Health.* 2011;5(4):117–24.
- Dasgupta A, Sarkar M. Menstrual hygiene: how hygienic is the adolescent girl? *Indian J Community Med.* 2008;33(2):77–80.
- Goel MK, Kundan M. Psycho-social behavior of urban Indian adolescent girls during menstruation. *Australas Med J.* 2011;4(1):49–52.
- Shanbhag D, Shilpa R, D'Souza N, Josephine P, Singh J, Goud BR. Perceptions regarding menstruation and Practices during menstrual cycles among high school going adolescent girls in resource-limited settings around Bangalore City, Karnataka, India. *Int J Collab Res Inter Med Public Health.* 2012;4(7):1353–62.
- Tiwari H, Oza UN, Tiwari R. Knowledge, attitudes and beliefs about Menarche of adolescent girls in Anand District, Gujarat. *East Mediterr Health J.* 2006;12(3-4):428–33.
- Thakre SB, Thakre SS, Reddy M, Rathi N, Pathak K, Ughade S. Menstrual hygiene: knowledge and practice among adolescent school girls of Saoner, Nagpur district. *J Clin Diagn Res.* 2011;5(5):1027–33.
- Yasmin S, Mallik S, Manna N, Ahmed A, Paria B. Menstrual hygiene among adolescent school students: an in-depth cross-sectional study in an urban community of West Bengal, India. *Sudan J Public Health.* 2013;8(2):60–4.
- Oche MO, Umar AS, Gana GJ et al.. Menstrual health: Unmet needs of adolescent girls in Sokoto, Nigeria. *Sci Res Essays.* 2012;7(3):410–8.
- Ray S, Dasgupta A. Determinants of menstrual hygiene among adolescent girls: a multivariate analysis. *Natl J Commun Med.* 2012;3(2):294–301.
- Boosey R, Prestwich G, Dave T. Menstrual hygiene management amongst Schoolgirls in the Rukungiri district of Uganda and their impact on their education: A cross-sectional study. *Pan African Med J.* 2014;19:253.
- Nemade D, Anjenaya S, Gujar R. Effect of health education on statistics and practices about menstruation among adolescent faculty girls of Kalamboli,

- Navi-Mumbai. fitness of Popul-Perspect issues. 2009;32(4):167-75
- Narayan okay, Srinivasa D, Pelto P, Veerammal S. Puberty Rituals, Reproductive Understanding, and Health of Adolescent Girls in South India. *Asia-percent Popul J.* 2001;16(2):225-38.
- ARORA A, Mittal A, Pathania D, Mehta C, Bungler R. Impact of health education on understanding and practices about menstruation among adolescent college women in the rural part of the district Ambala, Haryana. *Ind J Comm health.* 2013;25(4):492-7.
- Lawan UM, Yusuf NW, Musa AB. Menstruation and menstrual hygiene among adolescent college women in Kano, Northwestern Nigeria. *Afr J Reprod fitness.* 2010;14(3):201-7.
- Zegeye DT, Megabiaw B, Mulu A. Age at menarche and the menstrual pattern in younger secondary college humans in Northwest Ethiopia. *BMC Women's Fitness.* 2009;nine:29.
- Thakre SB, et al. Town-rural versions of menstrual troubles and practices of Female college students in Nagpur, India. *Indian Pediatr.* 2012;49(9):733-6.
- Udgiri R, Angadi MM, Patil S et al.. Expertise and practices concerning menstruation among adolescent women in a town slum in Bijapur. *J Indian Med Assoc.* 2010;108(8):514-6.
- Marvan ML, Molina-Abolnik M. Mexican youngsters' revel in of menarche and attitudes toward menstruation: function of communique among moms and daughters. *J Pediatr Adolesc Gynaecol.* 2012;25(6):358-63.
- Sharma M, Gupta S. Menstrual sample and abnormalities in the immoderate college girls of Dharan: A cross-sectional test of the boarding faculty of Nepal Med Coll J.2003;5(1):34-6.
- Adinma ED, Adinma JI. Menstrual traits among south-eastern Nigerian adolescent faculty women *West Afr J Med.* 2009;28(2):110-3.
- Reis N, Kilic D, Engin R, Karabulutlu O. Sexual and reproductive health desires of adolescent girls from conservative and low-income households in Erzurum, Turkey. *fitness of Popul Perspect trouble.* 2011;3(6):370-7.
- Bosch AM, Hutter I, van Ginneken JK. Perceptions of teens and their months for reproductive and sexual development in MATLAB, Bangladesh. *Int J Adolesc Med health.* 2008;20(three):329-42
- Dhingra R, Kumar A, Kour M. Understanding and Practices Associated with Menstruation Among Tribal (Gujjar) Adolescent Women. *Ethno-remedy.* 2009;3(1): 43-8
- El-Gilany AH. Badawi. El-Fedawy S. Menstrual hygiene among adolescent schoolgirls in Mansoura, Egypt. *Am. Reprod health subjects.* 2005;13(26):147-52.
- Gumanga SK, Kwame-Aryee RA. Menstrual trends in a few adolescents women in Accra, Ghana. *Ghana Med J.* 2012;46(1):3-7.
- Dambhare DG, Wagh SV, Dudhe JY. Age at menarche and menstrual cycle the patterns among adolescent women in India. *Glob J Health Sci.* 2012;4(1): a hundred and 5-11.
- Kotecha PV, et al. Reproductive fitness focuses on rural college-going young people in the Vadodara district. *Indian J sex Transm Dis.* 2009;30(2): 94-9.

- Mudey A, Kesharwani N, Mudey GA et al.. Pass-sectional observed attention concerning secure and hygienic practices among faculty-going adolescent girls in a rural area of Wardha District, India. *Glob J Health Sci.* 2010;2(2):225-31
- Ray S, et al. Knowledge and information on psychological, physiological, and gynecological problems among adolescent girls in eastern India. *Ethiopia J Health Sci.* 2011;21(3):183-9.
- Jarrah SS, Kamel AA. Attitudes and practices of school-aged girls towards menstruation. *Int J Nurs Pract.* 2012;18(3):308-15.
- Lee LK, et al. Menstruation among adolescent girls in Malaysia: A cross-sectional school survey. *Singapore Med J.* 2006;47(10):869-74.
- Wong LP. Attitudes toward menstruation, menstrual-related symptoms, and pre-menstrual syndrome among adolescent girls: A rural school-based survey. *Women's Health.* 2011;51(4):340-64.
- Wong LP. Premenstrual syndrome and dysmenorrhea: urban-rural and multipath differences in perception, impact, and treatment-seeking. *J Pediatr Adolesc Gynaecol.* 2011;24(5):272-7.
- Aniebue UU, Aniebue PN, Nwankwo TO. Impact of pre-menarcheal training on menstrual practices and hygiene in Nigerian schoolgirls. *Pan Afr Med J.* 2009;2:9.
- Iliyasu Z, et al. Sexual and reproductive health communication between mothers and adolescent daughters in Northern Nigeria. *Health Care Women Int.* 2012;33(2):138-52.
- Ajah LO, et al. Adolescent reproductive health challenges among schoolgirls in southeast Nigeria: Knowledge of menstrual patterns and contraceptive adherence. *Patient Preference Adherence.* 2015;9:1219-24.
- Chandraratne NK, Gunawardena NS. Premenstrual syndrome: The experience of a sample of Sri Lankan adolescents. *J Pediatr Adolesc Gynecol.* 2011;24(5):304-10.
- Abd El-Hameed NA, Mohamed MS, Ahmed NH, Ahmed ER. Assessment of dysmenorrhea and menstrual hygiene practices among adolescent girls in some nursing schools in LL-Minia governorate, Egypt. *J Am Sci.* 2011;7(9):216-23.
- Eswi A, Helal H, Elarousy W. Menstrual attitudes and knowledge of Egyptian female adolescents. *J Am Sci.* 2012;8(6):555-65.
- Omidvar S, Begum K. Factors influencing hygienic practices during menses among girls from South India: A cross-sectional study. *Int J Collab Res Intern Med Public Health.* 2010;2(12):411-23.
- Wong LP. Attitudes towards dysmenorrhea, impact, and treatment-seeking among adolescent girls: A rural school-based survey. *Aust J Rural Health.* 2011;19(4):218-23.
- Wong LP, Khoo EM. Menstrual-related attitudes and symptoms among Multiracial Asian adolescent females. *Int J Behav Med.* 2011;18(3):246-53.
- Sommer M. Ideologies of sexuality, menstruation, and risk: girls' experiences of puberty and schooling in northern Tanzania. *Cult Health Sex.* 2009;11(4):383-98.

- Crichton J, et al. Emotional and psychosocial aspects of menstrual poverty in resource-poor settings: A qualitative study of the experiences of adolescent girls in an informal settlement in Nairobi. *Health Care Women Int*. 2013;34(10):891-916.
- Mason L, et al. 'We keep it secret so no one should know' –a qualitative study to explore young schoolgirls' attitudes and experiences with menstruation in rural western Kenya. *PLoS One*. 2013;8(11):e79132.
- Munthali AC, Zulu EM. The timing and position of initiation rites in preparing younger human beings for formative years and accountable reproductive behavior in Malawi. *Afr J Reprod fitness*. 2007;11(three): hundred and 50-67.
- fifty-three. McMahon SA, et al. 'The girl together with her duration is the one to hang her head' Reflections on menstrual management amongst schoolgirls in rural Kenya. *BMC Int fitness haul rights*. 2011;eleven:7.
- Sommer M. An early window of possibility for promoting girls' health: Policy implications of the woman's puberty e-book task in Tanzania. *Int. Electron J Health Microbiol*. 2011; 14:77-92
- Dorgbetor G. Mainstreaming MHM in colleges through the play-primarily based approach: training discovered in Ghana. *Waterlines*. 2015;34(1): 41-50.56.
- Marvan ML, Vacio A, Espinosa-Hernandez G. Menstrual-associated changes expected with the aid of premenarcheal girls dwelling in rural and urban areas of Mexico. *Soc Sci Med*. 2003;56(4):863-8.
- Marvan ML, Vacio A, Espinosa-Hernandez G. A contrast of menstrual adjustments anticipated through pre-menarcheal kids and changes skilled with the aid of publish-menarcheal children in Mexico. *J Sch health*. 2001;71(9):458-61
- Pitangui AC, et al. Menstruation disturbances: incidence, characteristics, and effects on the daily activities of adolescent girls residing in Brazil. *J Pediatr Adolesc Gynecol*. 2013;26(three):148-52
- Santina T, Wehbe N, Ziade F. Exploring dysmenorrhea and menstrual reviews among Lebanese lady young people. *East Mediterr Health J*. 2012;18(8):857-63.
- Chaudhuri A, Singh A. How do school women cope with dysmenorrhea? *J Indian Med Assoc*. 2012; 10(5):287-91.
- Sommer M. Where the training machine and Girls' bodies collide: The Social and fitness impact of ladies' stories of menstruation and training in Tanzania. *J Adolesc*. 2010;33(4):521-9.
- Patil MS, Angadi MM. Menstrual patterns among adolescent girls in the rural regions of Bijapur. *Al Ameen J Med Sci*. 2013;6(1):17-20.
- Rana B, Prajapati A, Sonaliya KN, Shah V, Patel M, Solanki A. Assessment of menstrual hygiene practices among adolescent females in the Kheda district of Gujarat Kingdom, India. *Healthline J*. 2015;6(1):23-9.
- Sharma P, et al. Troubles associated with menstruation among adolescent girls. *Indian J Pediatr*. 2008; seventy-five (2): one hundred twenty-five-9, 65. Juyal R, Kandpal SD, Semwal J. Social elements of menstruation-associated practices in adolescent women in the district Dehradun. *Indian J*

- Network Fitness. 2013;25(three):213-6.
- Haque SE, et al. The impact of a school-based instructional intervention on menstrual health: An intervention examine among adolescent women in Bangladesh. *BMJ Open*. 2014;4(7):e004607.
- Bodat S, Ghate MM, Majumdar JR. School absenteeism during menstruation among rural adolescent girls in Pune. *Natl J Community Med*. 2013; four(2):212-6.
- Joshi D, Buit G, González-Botero D. Menstrual hygiene control: training and empowerment for women? *Waterlines*. 2015;34(1): 51-67.
- Sir Bernard Law et al. Sanitary pad interventions for girls' schooling in Ghana: A pilot study. *PLoS One*. 2012;7(10):e48274
- Oster E, Thornton R. Menstruation, sanitary products, and school attendance: Evidence from a randomized evaluation. *Am Econ J*. 2011;3(1):91-100.
- Mason L, Laserson K, Oruko K et al. Adolescent schoolgirls' experiences of Menstrual cups and pads in rural western Kenya: A qualitative study. *Waterlines*. 2015;34(1):15-30.
- Kabir H, et al. Treatment-seeking for selected reproductive health problems: Behaviors of unmarried female adolescents in two low-performing areas of Bangladesh. *Reprod Health*. 2014;11:54.
- Nair MK, et al. Menstrual disorders and menstrual hygiene practices of girls in higher secondary schools. *Indian J Pediatr*. 2012;79 Suppl 1:S74-8.
- Baidya S, Debnath M, Das R. Reproductive health problems among rural adolescent girls of the Mohanpur Block of the West Tripura District. *Al Ameen J Med Sci*. 2014;7(1):78-82.
- Wong LP, Khoo EM. Dysmenorrhea in a multiethnic population of adolescent Asian girls. *Int J Gynaecol Obstet*. 2010;108(2):139-42.
- Poureslami M. Assessing knowledge, attitudes, and behavior of adolescent girls in suburban districts of Tehran about dysmenorrhea and menstrual hygiene. *J Int Womens Stud*. 2002;3(2):51-61.
- Eryilmaz G, Ozdemir F. Evaluation of menstrual pain management approaches by Northeastern Anatolian adolescents. *Pain Manag Nurs*. 2009;10(1):40-7.
- Wasnik VR, Dhumale D, Jawarkar AK. A study of the menstrual pattern and problems among rural school-going adolescent girls in the Amravati district of Maharashtra, India. *Int J Res Med Sci*. 2015;33(55):1252-6.
- Fakhri M, et al. Promoting menstrual health among Persian adolescent girls from a low socioeconomic background: A quasi-experimental study. *BMC Public Health*. 2012;12:193.
- Allah ESA, Elsabagh EEM. Impact of a Health education intervention on Knowledge and Practice about Menstruation among female secondary school students in Zagazig City. *J Am Sci*. 2011;7(9):737-47.
- Sumpter C, Torondel B. A systematic review of the health and social effects of menstrual hygiene management. *PLoS One*. 2013;8(4):e62004.
- Nanda PMA, Mukherjee S, Barua A Mehl GL, Venkatraman CM. A study To evaluate the effectiveness of WHO tools: an orientation program on adolescent health for healthcare providers and adolescent job aid in India. Geneva: International Center for Research on Women, 2012.

- Vandenhoudt H, et al. Evaluation of a U.S. evidence-based parenting intervention in rural Western Kenya: From parents' matters! To families matter! *AIDS Educ Prev.* 2010;22(4):328–43.
- Sommer M, Ackatia-Armah N, Connolly S, Smiles D. A comparison of menstruation and education experiences of girls in Tanzania, Ghana, Cambodia, and Ethiopia. *Compare.* 2014;45(4):589–609.
- Children, S.t. Adolescent Sexual and Reproductive Health. 2015. Available from: http://www.savethechildren.org/site/c.8rKLIXMGIpI4E/b.9080949/k.F576/Adolescent_Sexual_and_Reproductive_Health.htm
- Health, I.f.R. Meeting the Needs of Adolescents: Introducing CCycle-Smart2013. Available from: <http://irh.org/blog/meeting-the-needs-of-adolescents-introducing-the-cycle-smart-kit/>
- Health, I.f.R. A3 project. Available from: http://irh.org/projects/a3_project/. Accessed 15 Oct 2014.
- Kettaneh APS, Todesco M. Good policy and practice booklet no. 9: puberty education and menstrual hygiene management. Paris: United Nations Educational, Scientific, and Cultural Organization, 2014.
- Always. Tips and advice: "The talk." Available from: <http://always.com/en-us/tips-and-advice/the-talk>. Accessed 15 Oct 2014.
- George R. Celebrating womanhood: How is better menstrual hygiene management the path to better health, dignity, and business? Geneva: Water Supply and Sanitation Collaboration Council, 2013.
- Sommer M. V. E., Worthington, N., Sahin M. WASH in schools empowers girl's education: proceedings of the menstrual hygiene management in schools virtual conference 2012. in *Menstrual Hygiene Management in Schools Virtual Conference*. New York, NY: United Nations Children's Fund and Columbia University; 2012.
- Kanotra SK, Bangal VB, Bhavthankar DP. Menstrual Patterns and Problems among adolescent girls in rural areas. *International Journal of Biomedical and Advance Research.* 2013; 4(8):551–